

Pharmacy informatics: A reform at drug consumption management in healthcare

Mahtab Karami ^{1*}

¹Department of Health Information Technology and Management, Health Information Management Research Center (HIMRC); School of Allied-Medical Sciences, Kashan University of Medical Sciences, Kashan, Iran

**Corresponding author: Department of Health Information Technology and Management, Health Information Management Research Center (HIMRC); School of Allied-Medical Sciences, Kashan University of Medical Sciences, Kashan, Iran. Email: mkarami33@gmail.com*

DOI: 10.22034/HBB.2017.23

Received: October 2, 2017; **Accepted:** October 30, 2017

In recent decades, since the pharmacy has a key role in patient care, advancement in information technology has created many opportunities for pharmacists and pharmacies to leverage informatics to support their practice and improve care [1].

Pharmacy informatics (PI) is an umbrella term that describes the use and integration of data, information, knowledge, and technology involved with medication use processes to improve outcomes. The use of informatics has ranged from improving pharmaceutical care in oncology to providing clinical decision support for antimicrobial stewardship and

pharmacokinetics to containing costs in managed care [2].

Generally, PI has two important applications including developments electronic prescribing systems and pharmacy information management systems. The first systems allow direct recording medication and treatment data in a computer system by physicians. The latter system collects, stores and manages the information related to the medications and their use in the patient care process [3, 4].

Application areas of PI are based on compliance with seven principles: "right patient", "right dose", "right route", "right drug", "right time",

"right information" and "proper documentation". Accordingly, these areas can be divided into four categories:

1. Prescribing management: Includes analysis of medication orders, control of drug-drug interactions by monitoring drug-drug, drug-food, and drug laboratory values incompatible, drug allergies and dosage according to age, weight, and other factors affecting patient which provide alert reminders for inconsistencies.
2. Drug distribution management: Includes demographic data, pharmaceutical therapeutic classification, drug identification number, generic and brand name, strength and dosage, drug manufacturer, date of distribution, and the amount of drug distribution.
3. Inventory Management: Includes maintenance integral index of inventories and pharmaceutical barcode to control previous and current medications, expiration date of drugs, control drug inventory by type, date and distributors, report income depot, and control of monetary statistics on purchased and delivered drugs as well as control of delivered drugs to patients with specific diseases.
4. Document Management: Includes the provision of alerts about the absence of any element of the documentation such as date, signature, and the name of receiving and delivering clerk [5-8].

For transition from traditional systems in the hospitals to the pharmaceutical informatics, it is necessary that these systems interact with other information systems in the hospital to support the pharmacy workflows. These support systems are as follows:

- Patient management systems that can provide patient's demographic information for the pharmacy information system (PIS).
- Electronic prescribing systems that feed and control the process of medication orders for PIS electronically in different parts of the hospital.
- Pathology information systems that can provide PIS to receive and display the results of laboratory tests related to certain drugs (for example: determining the plasma level for gentamicin or level of serum potassium for diuretics)
- Specific prescribing systems that can provide PIS to receive specific prescription such as chemotherapy
- Pharmacy robots that can be considered by means of labeling.
- Electronic cabinet in this way, the pharmacy computers are connected directly to the physicians and the pharmacists can remotely control drug cabinets in doctor office and after receiving the prescription; they can send signals to drug cabinets in remote to open a specific part of it. These cabinets contain barcoded drugs. The software sends the prescription to the patient

database to monitor the drug interactions and allergies, and then in the absence of danger for patient, it can confirm the medication administration. Additionally, patients can consult with pharmacist remotely [4, 9-11].

Using PI requires the infrastructures which support it effectively. These infrastructures can name the hardware that several users can simultaneously use it and change the locations. Another is wide bandwidth network to provide uninterrupted service at countrywide. Also, the data exchange based software to exchange the data easily between the different nodes or systems. The features of mobility and interoperability will also allow increased access to Telepharmacy.

On the other hand, both hardware and software systems need to be highly secured and protected against damage or loss of data. Data access levels can also determine who can access the information? How can this information be used? And finally, since healthcare services are based on social justice, access to medications and pharmacy in remote area is critical. To achieve this goal, the pharmacy information management system must be equipped with geographic information system [1-3, 5].

Using of PI reduces medication errors due to monitoring medication interactions, adverse effects, and allergies which can eliminate errors related to prescription handwriting.

Interoperability and data exchange between different systems such as patient management systems, pathology information systems, and specialized clinical applications may improve quality of care and patient safety in the organization and also can integrate healthcare services as well. Moreover, optimal management of resource consumption leads to cost saving and reduce healthcare costs too. Accordingly, it can be said that the use of PI in the health system is a revolution in drug consumption management [1, 7-9].

REFERENCES

- [1]. Baysari MT, Lehnbohm EC, Li L, Hargreaves A, Day RO, Westbrook JI. The effectiveness of information technology to improve antimicrobial prescribing in hospitals: A systematic review and meta-analysis. *Int J Med Inform*, 2016; 92: 15-34.
- [2]. Fox BI, Flynn AJ, Fortier CR, Clauson KA. Knowledge, skills, and resources for pharmacy informatics education. *Am J Pharm Educ*, 2011; 75(5): 93.
- [3]. Barnett J, Jennings H. Pharmacy information systems in Canada. *Student Health Technol Inform*, 2009; 143: 131-35.
- [4]. Patel J, Ogletree R, Sutterfield A, Pace JC, Lahr L. Optimized Computerized Order Entry Can Reduce Errors in Electronic Prescriptions

and Associated Pharmacy Calls to Clarify (CTC). *Appl Clin Inform*, 2016; 7(2): 587-95.

[5]. Farre A, Bem D, Heath G, Shaw K, Cummins C. Perceptions and experiences of the implementation, management, use and optimisation of electronic prescribing systems in hospital settings: protocol for a systematic review of qualitative studies. *BMJ Open*, 2016; 6(7): 1-6.

[6]. Niedrig D, Krattinger R, Jödicke A, Gött C, Bucklar G, Russmann S. Development, implementation and outcome analysis of semi-automated alerts for metformin dose adjustment in hospitalized patients with renal impairment. *Pharmacoepidemiol Drug Saf*, 2016; 25(10): 1204-09.

[7]. Vernaz N, Rollason V, Adlere L, Combescure C, Poncet A, Bonnabry P, Desmeules J. Snapshot of the prescribing practice for the clopidogrel and esomeprazole coprescription and cost evaluation of the application guidelines. *Pharmacol Res Perspect*, 2016; 4(3): 1-9.

[8]. Cresswell K, Mozaffar H, Shah S, Sheikh A. A systematic assessment of review to promoting the appropriate use of antibiotics through hospital electronic prescribing systems. *Int J Pharm Pract*, 2017; 25(1): 5-17.

[9]. Odukoya OK, Stone JA, Chui MA. E-prescribing errors in community pharmacies:

exploring consequences and contributing factors. *Int J Med Inform*, 2014; 83(6): 427-37.

[10]. Brenn BR, Kim MA, Hilmas E. Development of a computerized monitoring program to identify narcotic diversion in a pediatric anesthesia practice. *Am J Health Syst Pharm*, 2015; 72(16): 1365-72.

[11]. Tong V, Raynor DK, Aslani P. User testing as a method for identifying how consumers say they would act on information related to over-the-counter medicines. *Res Social Adm Pharm*, 2016; 7411(16): 30102-04.